



YAMASA CORPORATION

DIAGNOSTICS DEPARTMENT

Tokyo Office

1-23-8, Nihonbashi-kakigaracho

Chuo-ku, Tokyo, 103-0014, Japan

Tel. +81-3-3668-8558 Fax. +81-3-3668-8407

www.yamasa.com/shindan/index.htm

Monoclonal Antibody to Mouse β Interferon (Purified)

Code No. : 7891
Quantity: 500 μ L (1 mg)
Clone: 7F-D3
Quality: ACI rat IgG1 (Purified by Protein G chromatography.)
Solution: 2 mg/mL IgG, PBS, pH 7.4

Character:		Reactivity
to	Mouse IFN β	
	(L cell)	+
	(Recombinant)	+
	α, γ	-
	Human IFN	-

Example: Plate 3×10^4 cells/100 μ L of L cells in MEM containing 10% calf serum to a well of a micro plate.

Incubate at 37 °C for 24 hours.

Add 50 μ L of diluted 7891 and 50 μ L of mouse IFN- β in MEM.

Incubate at 37 °C for 24 hours.

Remove the culture medium.

Add 100 μ L of VSV in MEM.

Incubate at 37 °C for 24 hours.

Remove the culture medium.

Add 50 μ L of 0.05% crystal violet, 10% formalin, 10% ethanol.

Remove the culture medium.

Wash with tap water.

Dry the plate.

Score the CPE.

*At a concentration of 3.6 ng/ml, this antibody neutralizes 10 U/ml of mouse IFN- β to 1 U/ml.

Storage: The solution should be stored below -20 °C.

Repeated freezing and thawing should be avoided.

No preservative substance is used in this product.

Note: Research use only.

Reference: Kawade, Y. J. Interferon Res. 1, 61-70 (1980)