



YAMASA CORPORATION

DIAGNOSTICS DEPARTMENT

Tokyo Office

1-23-8, Nihonbashi-kakigaraicho


Chuo-ku, Tokyo, 103-0014, Japan

Tel. +81-3-3668-8558 Fax. +81-3-3668-8407

www.yamasa.com/shindan/index.htm

## Peroxidase Conjugated Monoclonal Antibody to Mouse Immunoglobulin Kappa Light Chain

Code No.: 7962  
 Quantity: 200  $\mu$ L (200  $\mu$ g)  
 Clone: 2B6  
 Quality: Horseradish peroxidase conjugated Fisher Rat IgG1.  
 Solution: 1 mg/mL Conjugate, 1% BSA, 0.1% ProClin 300 in PBS, pH7.4  
 Character:



**Table. Specificities of rat anti-mouse immunoglobulin antibodies**

Clone	Code. Purified/HRP Ab	Test clone <sup>a</sup> / isotype								
		1F4	2B11	1D1	5B11	3H8	2G11	2F2	2A10	1D8
		IgG1 $\kappa$	IgG1 $\kappa$	IgG1 $\lambda$	IgG1 $\lambda$	IgG2a $\kappa$	IgG2a $\kappa$	IgG2b $\kappa$	IgG2b $\kappa$	IgG3 $\kappa$
14G11	7965/7966	+	+	+	+	-	-	-	-	-
7D5	7967/7968	-	-	-	-	+	+	-	-	-
2B6	7961/7962	+	+	-	-	+	+	+	+	+
7G12	7963/7964	-	-	+	+	-	-	-	-	-

<sup>a</sup> Isotype identified mouse monoclonal antibodies were obtained from Yamasa monoclonal antibody libraries. Anti-mouse immunoglobulin antibodies were applied to mouse monoclonal antibodies-coated ELISA plates. Rat anti-mouse immunoglobulin antibodies that reacted the antigens were probed by horseradish peroxidase-labeled anti-rat IgG and TMB, a peroxidase substrate.  
 + absorbance > 1.2; - absorbance < 0.15.

**Storage:** The solution should be stored below -20°C.  
 Repeated freezing and thawing should be avoided.

**Note:** Research use only.