



YAMASA CORPORATION

DIAGNOSTICS DIVISION
Tokyo Office
1-23-8, Nihonbashi-kakigaracho
Chuo-ku, Tokyo, 103-0014, Japan
Tel.+81-3-3668-8558 Fax.+81-3-3668-8407
<https://diagnostics.yamasa.com/>

Anti-Human Interferon β Monoclonal Antibody

Code No. : 7889
Quantity: 500 μ L (1 mg)
Clone: B-02
Quality: BALB/c IgG1 κ (Purified by Protein A chromatography.)
Solution: 2 mg/mL IgG, PBS, pH 7.4

Character:		Reactivity
to Human IFN β	(Fibroblast)	++
	(Recombinant)	+
	α	-
	γ	-

Example:

1. Plate 3×10^4 cells/100 μ L of FL cells in MEM containing 10% calf serum to a well of a micro plate.
2. Incubate at 37 °C for 24 hours.
3. Add 50 μ L of diluted 7889 and 50 μ L/10 units of human fibroblast IFN- β in MEM.
4. Incubate at 37 °C for 24 hours.
5. Remove the culture medium.
6. Add 100 μ L of VSV in MEM.
7. Incubate at 37 °C for 24 hours.
8. Remove the culture medium.
9. Add 50 μ L of 0.05% crystal violet, 10% formalin, 10% ethanol.
10. Remove the culture medium.
11. Wash with tap water.
12. Dry the plate.
13. Score the CPE.

*At a concentration of 1.3 μ g/mL, this antibody neutralizes 10 U/mL of human IFN- β to 1 U/mL.

Storage: The solution should be stored below -20 °C.
Repeated freezing and thawing should be avoided.
No preservative substance is used in this product.

Note: Research use only.

Reference: Kawade, Y. J. Interferon Res. 1, 61-70 (1980)